

Asthma Action Card



Asthma and Allergy
Foundation of America

Reviewed by
 National Asthma Education and
Prevention Program



Name: _____ Grade: _____ Age: _____

Homeroom Teacher: _____ Room: _____

Parent/Guardian Name: _____ Ph: (H) _____

Address: _____ Ph: (W) _____

Parent/Guardian Name: _____ Ph: (H) _____

Address: _____ Ph: (W) _____

Emergency Phone Contact #1: _____
Name Relationship Phone

Emergency Phone Contact #2: _____
Name Relationship Phone

Physician Treating Student for Asthma: _____ Ph: _____

Other Physician: _____ Ph: _____

Emergency Plan

Emergency action is necessary when the student has symptoms such as _____,
_____, _____, _____,
_____, or has a peak flow reading of _____.

Steps to take during an asthma episode:

1. Check peak flow.
2. Give medications as listed below. Student should respond to treatment in 15-20 minutes.
3. Contact parent/guardian if: _____
4. Re-check peak flow.
5. Seek emergency medical care if the student has any of the following:
 - ✓ Coughs constantly
 - ✓ No improvement 15-20 minutes after initial treatment with medication and a relative cannot be reached
 - ✓ Peak flow of _____
 - ✓ Hard time breathing with:
 - Chest and neck pulled in with breathing
 - Stooped body posture
 - Struggling or gasping
 - ✓ Trouble walking or talking
 - ✓ Stops playing and can't start activity again
 - ✓ Lips or fingernails are grey or blue

ID Photo

Emergency Asthma Medications

Name	Amount	When to Use
1. _____		
2. _____		
3. _____		
4. _____		